



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 5/10, A01K 67/027, C07K 14/705, A61K 49/00	A1	(11) International Publication Number: WO 99/53032 (43) International Publication Date: 21 October 1999 (21.10.99)
(21) International Application Number: PCT/US99/08285 (22) International Filing Date: 15 April 1999 (15.04.99) (30) Priority Data: 60/115,516 15 April 1998 (15.04.98) US (71) Applicant (for all designated States except US): COR THERAPEUTICS, INC. [US/US]; 256 Grand Avenue, South San Francisco, CA 94080 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): LAW, Deborah, Ann [US/US]; 1428 5th Avenue, San Francisco, CA 94122 (US). PHILLIPS, David, R. [US/US]; 245 Parrot Drive, San Mateo, CA 94402 (US). (74) Agent: ADLER, Reid, G.; Morgan, Lewis & Bockius LLP, 1800 M Street, N.W., Washington, DC 20036 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: TRANSGENIC MAMMALS EXPRESSING MUTANT GP IIIa		
(57) Abstract <p>The present invention relates to mammals into which foreign DNA has been introduced or in which various modifications or substitutions have been made to an integrin β subunit, thereby generating transgenic or genetically-engineered non-human mammals. In particular, the present invention provides a transgenic mammal in which the endogenous GP IIIa gene has been replaced with an altered or mutant GP IIIa gene in which one or all of the phosphorylatable cytoplasmic tyrosine residues have been replaced with non-tyrosine residues such as phenylalanine. Since the platelets in the blood of the resultant transgenic mammals expressing the altered or mutant GP IIIa cannot undergo tyrosine phosphorylation to the extent that it occurs in wild-type mammals, these genetically-engineered animals provide a critical tool for assessing the importance of the phosphorylation reaction for platelet function. The invention is also useful for studying the effect of the mutant GP IIIa integrin subunit on biological processes other than platelet formation.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TRANSGENIC MAMMALS EXPRESSING MUTANT GP IIIa

RELATED APPLICATIONS

This application is related to U.S. Patent Application No. 08/734,607, filed
5 October 18, 1996; U.S. Provisional Application No. 60/031,665, filed November 21, 1996;
U.S. Provisional Application No. 60/042,093, filed March 28, 1997; and, U.S. Patent
Application No. 08/975,653, filed November 21, 1997; all of which are hereby
incorporated by reference in their entireties.

10 FIELD OF THE INVENTION

The present invention relates to mammals into which foreign DNA has been
introduced or in which certain codons of integrin-encoding genes have been replaced,
thereby generating transgenic or genetically-engineered mammals. In particular, the
present invention provides a transgenic mammal in which the endogenous GP IIIa gene
15 (also known as $\beta 3$) has been replaced in whole or in part with a mutant GP IIIa gene in
which one or both of the two phosphorylatable cytoplasmic tyrosine residues have been
replaced with non-tyrosine residues, for example, such as phenylalanine. Since the cells,
platelets in particular, that are found in the blood of the resultant transgenic mammals that
express an altered GP IIIa gene cannot undergo tyrosine phosphorylation, either in whole
20 or in part, as occurs in their wild type counterparts, these animals provide a useful tool for
assessing the importance of the phosphorylation of these tyrosine residues for platelet
function. The present invention is also useful for studying the effect of the mutant GP IIIa
integrin subunit on biological processes other than platelet formation.

25 BACKGROUND OF THE INVENTION

All of the publications and patent applications that are identified in this
specification are hereby incorporated by reference to the same extent as if each individual
publication or patent application was specifically and individually indicated to be
incorporated by reference.

A. Integrins

Integrins are a family of $\alpha\beta$ heterodimers that mediate adhesion of cells to extracellular matrix proteins and to other cells (Clark *et al.*, Science (1995) 268:233-239). Integrins also bind to the actin cytoskeleton through a series of intermediate proteins, and thus provide a link between the extracellular matrix and the intracellular cytoskeleton and its associated motile machinery. Such transmembrane linkages are required for cell migration. Many biological responses are dependent at least to some extent upon integrin-mediated adhesion and cell migration, including embryonic development, hemostasis, clot retraction, mitosis, angiogenesis, inflammation, immune response, leukocyte homing and activation, phagocytosis, bone resorption, tumor growth and metastasis, atherosclerosis, restenosis and wound healing.

Members of the integrin family also participate in signal transduction. This is evidenced by an alteration in the adhesive affinity of cell surface integrins in response to cellular activation, termed inside-out signal transduction. Additionally, effects on intracellular signaling pathways following integrin-mediated adhesion have been observed, termed outside-in signal transduction.

The integrin family consists of 15 related known α subunits ($\alpha 1$, $\alpha 2$, $\alpha 3$, $\alpha 4$, $\alpha 5$, $\alpha 6$, $\alpha 7$, $\alpha 8$, $\alpha 9$, αE , αV , αIIb , αL , αM , and αX) and 8 related known β subunits ($\beta 1$, $\beta 2$, $\beta 3$, $\beta 4$, $\beta 5$, $\beta 6$, $\beta 7$, and $\beta 8$). Luscinskas *et al.*, *FASEB J.*, 8: 929-938 (1994). Integrin α and β subunits are known to exist in a variety of pairings as indicated in Figure 1. Integrin ligand specificity is determined by the specific pairing of the α and β subunits, although some redundancy exists as several of the integrins are known to bind the same ligand.

Two known pairings of the $\beta 3$ subunit have been observed: with αV to make $\alpha V\beta 3$, the Vitronectin Receptor; and with GP IIb to make GP IIb-IIIa, the Fibrinogen Receptor. $\alpha V\beta 3$ is widely distributed, is the most promiscuous member of the integrin family and mediates cellular attachment to a wide spectrum of adhesive proteins, mostly at the R-G-D sequence on the adhesive protein. The biological processes mediated by $\alpha V\beta 3$ are diverse and include bone resorption, angiogenesis, tumor metastasis and restenosis. $\alpha V\beta 3$ is known to signal upon adhesive protein ligation (P.I. Leavesley, *et al.*, *J. Cell Biol.* 121:163-170 (1993)). As an example, endothelial cells undergo apoptosis when relieved of ligation (P.C. Brooks, *Cell* 79:1157-1164 (1994)).

B. Interaction of Integrins with Known Cytoskeletal Proteins

The binding of unmodified α and β subunit cytoplasmic domains of integrins to a variety of cytoskeletal and signaling proteins has been documented. S. Dedhar *et al.*,
5 *Curr. Opin. Cell Biol.* 8:657-669 (1996). Morphological studies have shown that many of these proteins are concentrated in focal adhesions where integrins cluster and bind to both the extracellular matrix and cytoskeletal proteins. I. Knezevic *et al.*, *J. Biol. Chem.* 271(27):16416-16521 (1996).

For example, talin, a 235 kD vinculin and actin binding protein, binds to the
10 cytoplasmic domains of α IIb and β 3 in a solid phase binding assay. I. Knezevic *et al.*, *Id.* The binding of α actinin, a 100 kD vinculin binding protein and actin cross-linking protein, to the cytoplasmic domain of β 1 and β 3 in solid phase binding assays has also been observed. C.A. Otey *et al.*, *J. Biol. Chem.* 268(28):21193-21197 (1993); and C. A. Otey *et al.*, *J. Cell Biol.* 111:721-729 (1990). Binding studies have demonstrated an
15 interaction between the cytoplasmic domain of β 1 and tensin, a 215 kD SH2 domain containing vinculin and actin binding protein. S. Lin *et al.* *Mol. Biol. Cell* 7 Supp. 389a, Abstract 2259 (1996).

Other cytoskeletal related proteins also interact with integrins. Skelemin, a 195 kD myosin and intermediate filament binding protein, binds to the membrane proximal
20 regions of β 1 and β 3 cytoplasmic domains. K.B. Reddy *et al.*, *Mol. Biol. Cell* 7 Supp. 385A, Abstract 2237 (1995). These authors suggested that skelemin could link myosin and intermediate filaments to β integrins.

Paxillin, a vinculin binding signaling protein also binds to the cytoplasmic domain of the β 1 integrin. M.D. Schaller *et al.*, *J. Cell Biol.* 130:1181-1187 (1995). It is not yet
25 known whether the β 1-paxillin association is direct or indirect, however paxillin was postulated as being the substrate for and tyrosine phosphorylated by tyrosine kinase pp125 FAK. The actin binding protein filamin has been shown to bind to the cytoplasmic tail of the β 2 integrin subunit in vitro and co-immunoprecipitated and co-localized with β 2 integrins in vivo. C.D. Sharma *et al.*, *J. Immunol.* 154: 3461-3470 (1995).

Chain kinase family of serine/threonine kinases has also been reported. Walker *et al.*, *Mol.*

Biol. Cell 7 Supp. 385A, Abstract 2235 (1995). This kinase was said to be part of a complex of proteins including α -actinin and myosin, however, it was unclear whether the kinase associated directly with the cytoplasmic tails of integrins or through a complex of proteins.

Although the cytoskeletal proteins listed above have been shown to interact with cytoplasmic domains of integrin subunits with purified proteins or peptides, it is not known how these interactions occur within cells or how these interactions are regulated. Furthermore, the integrin/cytoskeletal interactions described thus far do not occur in a phosphotyrosine-dependent manner.

C. Tyrosine Phosphorylation of the Cytoplasmic Domain of Integrin β Subunits

Platelet aggregation induced by a number of agonists results in the phosphorylation of tyrosine residues in the $\beta 3$ cytoplasmic tail. Law *et al.*, *J. Biol. Chem* 271:10811-10815 (1996). In some respects, the phosphorylation of both tyrosine residues was necessary for binding to certain signaling proteins, whereas other signaling proteins bound following monophosphorylation. Furthermore, adhesion to vitronectin by cells transfected with $\alpha v \beta 3$ induces a robust tyrosine phosphorylation of the $\beta 3$ subunit. Blystone *et al.*, *J. Biol. Chem* 271:31458-31462 (1996).

Studies have shown that the sequences of the cytoplasmic domains of $\beta 1$, $\beta 2$ and $\beta 3$ which contain tyrosines are important for normal integrin/cytoskeletal interactions. For example, the substitution of tyrosine 747 by alanine in $\beta 3$ transfected into CHO cells abolished $\beta 3$ -mediated cell spreading, blocked the recruitment of $\alpha IIb \beta 3$ to preestablished adhesion plaques, and decreased the ability of $\alpha IIb \beta 3$ to mediate internalization of fibrinogen-coated particles. J. Ylanne *et. al.*, *J. Biol. Chem.*, 270, 9550-9557, (1995).

Additional experiments reported by Ylanne *et al.*, *Id.*, showed further that substitution of alanine for tyrosine 759 decreased cell spreading and the recruitment of $\alpha IIb \beta 3$ to adhesion plaques, while deletion of the carboxy terminal pentapeptide that contains this sequence had an even more pronounced effect on the function of the integrin. These authors concluded integrin-mediated cell spreading does not occur because the factors that are absolutely required for integrin-mediated cell spreading cannot bind either

the $\beta 3$ truncated at residue 757 or the integrin with tyrosine 747 of $\beta 3$ substituted by alanine.

Point mutations in homologous domains in $\beta 1$ - and $\beta 2$ -containing integrins also modulate function, as these mutations affect integrin-cytoskeletal interactions by
5 reducing focal adhesions, A. A. Reszka et. al., *J. Cell Biol.* 117:1321-1330 (1992), and integrin activation, M. L. Hibbs et. al., *J. Exp. Med.* 174:1227-1238 (1991), respectively. Tyrosine kinases similarly were found to be essential in regulating the cytoskeletal attachment of $\alpha \text{IIb}\beta 3$. Schoenwaelder et al., *J. Biol. Chem.* 269(51):32479-32487 (1994).

Overall, the interactions between the two tyrosines and the "cell adhesion
10 regulatory domain" or "CARD" of residues 747-762 of the $\beta 3$ cytoplasmic domain were reported to be essential for regulation of the adhesive function of integrin $\beta 3$. Liu et al., *PNAS* 93:11819-11824 (1996). A 16-amino acid sequence from the CARD inhibited adhesion of HEL and ECV 304 cells to immobilized fibrinogen by competing with
intracellular protein-protein interactions that "engage the business end" of the integrin $\beta 3$
15 tail. However, the identity of cytoplasmic protein(s) interacting with CARD was said to remain to be established.

D. Myosin

The platelet plasma membrane is coated by a lattice-like structure, known as the
20 membrane skeleton, that is composed of short actin filaments, actin-binding protein, spectrin, vinculin and various other proteins, not all yet identified. Fox et al., *J. Biol. Chem.* 268(34):25973-25984 (1993). On the cytoplasmic side, the skeleton appears to be associated with a network of cytoplasmic actin filaments. The membrane skeleton coats the lipid bilayer and is associated with both extracellular glycoproteins and intracellular
25 cytoskeletal elements. Fox et al. suggested that GPIIb-IIIa induces redistribution of components of the membrane skeleton and associated signaling molecules as a step in regulating integrin-induced motile events in platelets.

Myosin is a contractile protein that interacts with actin to produce contraction or movement. The term "myosin" broadly refers to a diverse superfamily comprised of at

myosins is their ability to reversibly bind to actin and to hydrolyze MgATP. See Figure 5 and J. R. Sellers and H.V. Goodson, Protein Profile 2:1323-1339 (1995).

All types of myosin that have been purified are multimeric and appear to possess at least three functional domains- a head, neck and tail. The head or motor domain contains nucleotide and actin binding sites and is the most conserved region of the myosin superfamily. The neck domain consists of a long single alpha helical strand from the heavy chain which is stabilized by the binding of light chain subunits. The tail region, which serves to anchor myosin so that it can translocate actin, is the most diverse primary sequence of all the regions and may serve to anchor certain myosin isoforms to cell or organelle membranes. It has been suggested that myosin clustering within a cell may occur on membranes or on actin filaments themselves. Titus, *Trends in Cell Biology* 7:119 (1997). However, the precise biochemical mechanism of interaction between the myosin tail and cytoplasmic structures has not heretofore been described.

15 **E. Signal Transduction**

The involvement of the cytoplasmic domain of GP IIb-IIIa in integrin signal transduction is inferred from mutagenesis experiments. Deletion of the cytoplasmic domain of GP IIb results in a constitutively active receptor that binds fibrinogen with an affinity equivalent to the wild-type complex, implying that the cytoplasmic tail of GP IIb has a regulatory role (T. E. O'Toole, *et al.*, *Cell Regul.* 1:883-893, (1990)). Point mutations, deletions and other truncations of GP IIb-IIIa affects the ligand binding activity of GP IIb-IIIa and its signaling response (P. E. Hughes, *et al.*, *J. Biol. Chem.* 270:12411-12417, (1995); J. Ylanne, *et al.*, *J. Biol. Chem.* 270:9550-9557, (1995)).

Chimeric, transmembrane proteins containing the cytoplasmic domain of GP IIIa, but not of GP IIb, inhibit the function of GP IIb-IIIa (Y.P. Chen *et al.*, *J. Cell Biol.* 269:18307-18310, (1994)), implying that free GP IIIa cytoplasmic domains bind proteins within cells which are necessary for normal GP IIb-IIIa function. Several proteins have been shown to bind either the transmembrane domains or the cytoplasmic domains of GP IIb or GP IIIa.

CD-9, a member of the tetraspanin family of proteins (F. Lanza, *et al.*, *J. Biol. Chem.* 266:10638-10645, 1991), has been found to interact with GP IIb-IIIa on aggregated

platelets. $\beta 3$ -endonexin, a protein identified through two hybrid screening using the cytoplasmic domain of GP IIIa as the "bait", has been found to interact directly and selectively with the cytoplasmic tail of GP IIIa (S. Shattil *et al.*, *J. Cell. Biol.* 131:807-816, (1995)). $\beta 3$ -endonexin shows decreased binding to the GP IIIa cytoplasmic domain containing the thrombasthenic S752-P mutation. It is not yet known whether either of these GP IIIa-binding proteins are involved in signal transduction.

Cytoplasmic proteins that bind to $\alpha V\beta 3$ have also been described which may be interacting with the integrin at the GP IIIa cytoplasmic domain sequence. Bartfeld and coworkers (N. S. Bartfeld *et al.*, *J. Biol. Chem.* 268:17270-17276, (1993)) used immunoprecipitation from detergent lysates to show that a MW=190-kDa protein associates with the $\alpha V\beta 3$ integrin from PDGF-stimulated 3T3 cells. IRS-1 was found to bind to the $\alpha V\beta 3$ integrin following insulin stimulation of Rat-1 cells stably transfected with DNA encoding the human insulin receptor (K. Vuori and E. Ruoslahti, *Sci.* 266:1576-1578, (1994)). Kolanus *et al.* (*Cell* 86:233-242, (1996)) recently identified Cytohesin-1. Cytohesin-1 specifically binds to the intracellular portion of the integrin $\beta 2$ chain, and overexpression of cytohesin-1 induces $\beta 2$ integrin-dependent binding of Jurkat cells to ICAM-1. A novel serine/threonine kinase, ILK-1, was found to associate with the $\beta 1$ cytoplasmic domain (Hannigan *et al.*, *Nature* 379:91-96, (1996)). Overexpression of ILK-1 inhibits adhesion to the integrin ligands fibronectin, laminin, and vitronectin.

Integrin binding to adhesive proteins and integrin signal transduction have a wide variety of physiological roles, as identified above. Enhanced signaling through integrins allows for increased cell adhesion and activation of intracellular signaling molecules which causes enhanced cell mobility and growth, enhanced cell responsiveness, and modulations in morphological transformations. Although integrins responsible for cellular function have been described and signaling events are beginning to be elucidated, the mechanism by which integrins transduce signals remains to be determined. To understand the molecular mechanisms of the inside-out and outside-in signaling roles mediated by the cytoplasmic tails of $\beta 3$ integrin requires the identification of the intracellular molecules that interact with the intracellular tails of integrin. It has been reported that α -actinin binds

as a $\beta 1$ interacting protein but ILK-1 does not bind to $\beta 3$ (Hannigan *et al.*, *Nature* 379:91-96 (1996)).

F. Homologous Recombination

5 Genes can be introduced in a site directed fashion using homologous recombination. This can be used in the creation of a transgenic animal, wherein the animal would be mutated, and the phenotype of the mutation could be studied for purposes of drug screening, investigating physiologic processes, developing new products and the like. Papers discussing homologous recombination are discussed in R. Kucherlapati *et al.*,
10 (1995) U.S. Patent No. 5,413,923.

Homologous recombination permits site-specific modifications in endogenous genes and thus inherited or acquired mutations may be corrected, and/or novel alterations may be engineered into the genome. The application of homologous recombination to gene therapy depends on the ability to carry out homologous recombination or gene
15 targeting in normal, somatic cells for transplantation.

To prepare cells for homologous recombination, embryonic stem cells or a stem cell line may be obtained. Cells other than embryonic stem cells can be utilized (*e.g.* hematopoietic stem cells etc.) (See for more examples, J.G. Seidman *et al.*, (1994) U.S. Patent No. 5,589,369). The cells may be grown on an appropriate fibroblast fetal layer or
20 grown in the presence of leukemia inhibiting factor (LIF) and then used. The embryonic stem cells may be injected into a blastocyst, that has been previously obtained, to provide a chimeric animal. The main advantage of the embryonic stem cell technique is that the cells transfected with the "transgene" can be tested prior to reimplantation into a female animal for gestation for integration and the effect of the transgenes. In contrast to the
25 conventional microinjection technique, the homologous respective endogenous gene can be removed from a chromosome by homologous recombination with the transgene. By subsequent cross-breeding experiments, animals can be bred which carry the transgene on both chromosomes. If mutations are incorporated into the transgenes which block expression of the normal gene production, the endogenous genes can be eliminated by this
30 technique and functional studies can thus be performed.

Homologous recombination can also proceed extrachromasomally, which may be of benefit when handling large gene sequences (e.g., larger than 50 kb). Methods of performing extrachromosomal homologous recombination are described in R.M Kay *et al.*, (1998) U.S. Patent No. 5,721,367.

5

G. Production of Transgenic Animals

Transgenic animals are genetically modified animals into which cloned genetic material has been experimentally transferred. The cloned genetic material is often referred to as a transgene. The nucleic acid sequence of the transgene is integrated at a locus of a genome where that particular nucleic acid sequence is not otherwise normally found. The transgene may consist of nucleic acid sequences derived from the genome of the same species or of a different species than the species of the target animal.

The development of transgenic technology allows investigators to create mammals of virtually any genotype and to assess the consequences of introducing specific foreign nucleic acid sequences on the physiological and morphological characteristics of the transformed animals. The availability of transgenic animals permits cellular processes to be influenced and examined in a systematic and specific manner not achievable with most other test systems. For example, the development of transgenic animals provides biological and medical scientists with models that are useful in the study of disease. Such animals are also useful for the testing and development of new pharmaceutically active substances.

Transgenic animals can be produced by a variety of different methods including transfection, electroporation, microinjection, gene targeting in embryonic stem cells and recombinant viral and retroviral infection (*see, e.g.*, U.S. Patent No. 4,736,866; U.S. Patent No. 5,602,307; Mullins *et al.*, *Hypertension* 22(4):630-633 (1993); Brenin *et al.*, *Surg. Oncol.* 6(2)99-110 (1997); Tuan (ed.), *Recombinant Gene Expression Protocols*, Methods in Molecular Biology No. 62, Humana Press (1997)). The term "knock-out" generally refers to mutant organisms, usually mice, which contain a null allele of a specific gene. The term "knock-in" generally refers to mutant organisms, also usually mice, into which a

A number of recombinant murines have been produced, including those which express an activated oncogene sequence (U.S. Patent No. 4,736,866); express simian SV 40 T-antigen (U.S. Patent No. 5,728,915); lack the expression of interferon regulatory factor 1 (IRF-1) (U.S. Patent No. 5,731,490); exhibit dopaminergic dysfunction (U.S. Patent No. 5,723,719); express at least one human gene which participates in blood pressure control (U.S. Patent No. 5,731,489); display greater similarity to the conditions existing in naturally occurring Alzheimer's disease (U.S. Patent No. 5,720,936); have a reduced capacity to mediate cellular adhesion (U.S. Patent No. 5,602,307); possess an bovine growth hormone gene (Clutter et al., *Genetics* 143(4):1753-1760 (1996)); and, are capable of generating a fully human antibody response (McCarthy, *The Lancet* 349(9049):405 (1997)).

While murines, especially mice and rats, remain the animals of choice for most transgenic experimentation, in some instances it is preferable or even necessary to use alternative animal species. Transgenic procedures have been successfully utilized in a variety of non-murine animals, including sheep, goats, pigs, dogs, cats, monkeys, chimpanzees, hamsters, rabbits, cows and guinea pigs (see, e.g., Kim et al., *Mol. Reprod. Dev.* 46(4):515-526 (1997); Houdebine, *Reprod. Nutr. Dev.* 35(6):609-617 (1995); Petters, *Reprod. Fertil. Dev.* 6(5):643-645 (1994); Schniekc et al., *Science* 278(5346):2130-2133 (1997); and, Amoah, *J. Animal Science* 75(2):578-585 (1997)).

The method of introduction of nucleic acid fragments into recombination competent mammalian cells can be by any method which favors co-transformation of multiple nucleic acid molecules. Detailed procedures for producing transgenic animals are readily available to one skilled in the art, including the recitations in U.S. Patent No. 5,489,743 and U.S. Patent No. 5,602,307.

SUMMARY OF THE INVENTION

The present invention provides non-human mammals comprising altered, substituted or mutant integrin cytoplasmic β subunit genes (and their expression products) in which at least one of the two (or more) cytoplasmic tyrosine residues of the expression product are replaced with a non-tyrosine residue such as phenylalanine. Examples of

mammals encompassed by the present invention include sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

More specifically, the present invention provides non-human mammals comprising a mutant GP IIIa gene and expression product wherein one or both of the cytoplasmic tyrosine residues that are capable of phosphorylation have been replaced with a non-phosphorylatable residue such as phenylalanine.

Even more specifically, the present invention provides transgenic mice comprising a mutant GP IIIa gene wherein the two cytoplasmic tyrosine residues 747 and 759 have each been replaced with phenylalanine.

The present invention also provides platelets isolated from the blood plasma of the transgenic non-human mammals of the present invention.

The present invention also provides methods of preparing a transformed non-human mammal with a mutant integrin cytoplasmic β subunit gene, such as the GP IIIa gene, wherein, for example, at least one of the two tyrosine residues of the endogenous GP IIIa gene has been replaced with a non-tyrosine residue to prepare the mutant GP IIIa, wherein the methods comprise: a) introducing into embryonic stem cells a nucleic acid molecule encoding the mutant integrin cytoplasmic β subunit gene, such as a mutant or altered GP IIIa gene, and b) regenerating a transformed non-human mammal from the cells resulting from step a).

More specifically, the present invention provides methods of preparing a transformed non-human mammal with, for example, a mutant GP IIIa gene wherein at least one of the two tyrosine residues of the endogenous GP IIIa gene has been replaced with a non-tyrosine residue to prepare the mutant GP IIIa, said method comprising:

- a) introducing into embryonic stem cells a nucleic acid molecule encoding the mutant GP IIIa gene and a selectable marker flanked by FRT sites;
- b) identifying and selecting transformed cells;
- c) removing the selectable marker from the transformed cells selected in step b) by transient transformation with FLP recombinase;

- e) regenerating a transformed non-human mammal from the blastocysts of step d), wherein the regenerated transformed non-human mammal is chimeric for the mutant GP IIIa gene;
- f) breeding the chimeric non-human mammal with a wild-type non-human mammal to produce a non-human mammal heterozygotic for the mutant GP IIIa gene;
- g) crossing a heterozygotic non-human mammal produced in step f) with a second heterozygotic non-human mammal produced in step f) ; and,
- h) selecting a non-human mammal homozygotic for the mutant GP IIIa gene from the resulting progeny.

The present invention also provides methods of comparing a characteristic between two mammals of the same species, or strain, wherein one mammal has, for example, a wild-type GP IIIa gene and the other mammal has an altered or mutant GP IIIa gene, wherein at least one of the two tyrosine residues of the wild-type GP IIIa gene has been replaced with a non-tyrosine residue in the mutant GP IIIa gene. More specifically, the present invention includes such methods using transgenic mammals wherein both cytoplasmic tyrosine residues of GP IIIa have been replaced with a non-tyrosine residue such as phenylalanine. Even more specifically, the present invention includes such methods using transgenic mammals wherein the cytoplasmic tyrosine residues 747 and 759 have both been replaced with a non-tyrosine residue such as phenylalanine.

Examples of characteristics which can be compared use in the transgenic mammals of the present invention include comparing various physiological or biological functions and responses that are mediated, in whole or in part, by the cytoplasmic β subunit gene of various integrin molecules. For example, a comparison of such functions would include observations and comparisons of the bleeding or clotting time between the two mammal types (i.e., between wild-type and genetically engineered animals); comparing various thrombotic responses between the two mammal types; comparing the level, extent and rate of angiogenesis between the two mammal types in response to various angiogenic stimuli;

comparing the rate and extent of tumor metastasis between the two mammal types; and comparing scope, rate and extent of inflammation between the two mammal types.

The present invention also provides methods of determining the effect of various agents on selected biological characteristics of a genetically engineered mammal
5 expressing an altered or mutant integrin β subunit gene that is attributable to the expression of the GP IIIa gene, wherein the methods comprise: a) administering said agent to the transgenic mammal; b) maintaining said mammal for a desired period of time after said administration; and, c) determining whether a characteristic of said mammal that is attributable to the expression of the mutant GP IIIa gene has been affected by the
10 administration of said agent.

Further objects and advantages of the present invention will be clear from the description and examples which follow. For example, even though the examples are directed to transgenic animals comprising mutant $\beta 3$ subunits, the present invention can be practiced with other integrin β subunits that have one or more phosphorylatable tyrosine
15 residues in the endogenous gene coding for that particular integrin β subunit gene.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the pairing of α and β integrin subunits.

20 Figure 2 shows the cytoplasmic domain of various integrin subunits.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this
25 invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described.

I. General Description

FIG. 1 is a schematic diagram of the pairing of α and β integrin subunits. FIG. 2 is a schematic diagram of the cytoplasmic domain of various integrin subunits.

application were described by David R. Phillips at the American Heart Association meeting in San Francisco, California on April 15, 1998. The content of that presentation and any printed material related thereto are herein incorporated by reference in their entirety.

5 Previous work by the present inventors indicated that the tyrosine phosphorylation of the GP IIIa subunit of the platelet integrin GP IIb-IIIa (also known as α IIb β 3) could modulate signal transduction by this integrin (see, e.g., U.S. Patent Application No. 08/975,653, filed November 21, 1997, which is incorporated by reference herein). That GP IIIa tyrosine phosphorylation was dependent upon platelet aggregation, and the fact
10 that signaling proteins as well as cytoskeletal proteins became associated with the phosphorylated GP IIIa, suggested to us that GP IIIa phosphorylation was important in outside-in GP IIb-IIIa signaling in platelets. Successful GP IIb-IIIa-mediated outside-in signaling is required for platelet functions such as the formation of stable platelet aggregates, an important process for normal hemostasis and one which, under aberrant
15 conditions, can lead to the formation of occlusive thrombi. Thus, if tyrosine phosphorylation of GP IIIa is indeed a critical step in outside-in GP IIb-IIIa signaling, then it could be hypothesized that compounds capable of inhibiting such a phosphorylation event would have anti-thrombotic properties. To address the importance of tyrosine phosphorylation of GP IIIa in platelet function a mutant mouse has been generated in
20 which the endogenous GP IIIa gene was replaced with one in which the two cytoplasmic tyrosine residues were mutated to phenylalanines (GP IIIa (Y747F, Y759F)). The platelets from such animals express GP IIIa that can not undergo tyrosine phosphorylation and therefore provide a critical tool for assessing the importance of the phosphorylation reaction for platelet function.

25 Although assessing the effect of GP IIIa mutations in transfected cell lines (i.e., *in vitro*) is useful in determining the role of various residues in GP IIIa function, the importance of the GP IIIa cytoplasmic tyrosine residues in platelet function can best be determined *in vivo*. Thus we undertook a gene replacement project in which the endogenous murine GP IIIa gene was replaced by a mutant genes with at least one non-
30 tyrosine substitution at one of the two cytoplasmic tyrosine residues. More specifically, the endogenous murine GP IIIa gene was replaced by a mutant gene with double tyrosine

to phenylalanine substitutions in the cytoplasmic domain at positions 747 and 759 respectively. Since the murine GP IIIa gene has not been sequenced, P1 clones containing murine genomic GP IIIa were obtained using PCR-based technology with PCR primers based on the sequence of the human GP IIIa gene. DNA corresponding to the area of interest, *i.e.*, the two exons encoding the GP IIIa cytoplasmic domain, was mapped using common molecular biology techniques. The two tyrosines in the cytoplasmic domain (Y747 and Y759) were mutated to phenylalanines and a targeting vector was constructed which included the mutated GP IIIa genomic DNA as well as a selectable drug marker (neomycin resistance) flanked by FRT sites, the recognition sequences for the FLP recombina-
5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

recombinase. This vector was transfected into embryonic stem cells and cells which had undergone a homologous recombination event were identified by Southern blotting and by PCR using specific primers. Those ES cells that contained one normal GP IIIa allele and one GP IIIa (Y747F, Y759F) mutant allele as well as the neomycin resistance DNA were then transiently transfected with the FLP recombina-
5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

Again, ES cells that had undergone the desired event were identified by both Southern blotting and PCR. The now standard protocols for the generation of mutant mice were employed. Basically, the desired mutant ES cells being injected into blastocysts in order to generate chimeric mice which were bred to wild-type mice to produce heterozygote animals expressing one normal and one mutant GP IIIa allele (as assessed by Southern blotting of tail genomic DNA and PCR). These heterozygote animals were then crossed to generate litters in which ~1/4 of the animals were homozygote for mutant GP IIIa (*ie.* the only GP IIIa expressed by these mice contained the double tyrosine to phenylalanine mutation).

These transgenic mice provide a critical tool for assessing the importance of the GP IIIa cytoplasmic tyrosine residues on platelet function. Previous work indicated that phosphorylation of the GP IIIa tyrosine residues was important in outside-in GP IIb-IIIa signaling. The platelets from the mutant mice express a GP IIb-IIIa in which the GP IIIa can not be tyrosine phosphorylated. Thus platelets from these mice can be used in a number of assays to assess the role of outside-in GP IIb-IIIa signaling on platelet function

obligatory part of outside-in signaling, one would predict the mutant mice to be defective in thrombotic responses, where the formation of very large platelet aggregates is required. We can now employ murine thrombotic models with these mutant mice to directly assess this issue. Indeed, this mutant mouse will provide information on the utility of generating
5 therapeutics designed to interfere either directly with GP IIIa tyrosine phosphorylation, or with an event dependent on this phosphorylation e.g. interaction of phospho-GP IIIa with an adaptor protein.

Expression of the GP IIIa (β_3) integrin subunit is not restricted to platelets. In endothelial cells this protein pairs with the α_v subunit to form the $\alpha_v\beta_3$ integrin. This
10 integrin appears to play an important role in angiogenesis and tumor metastasis. In neutrophils, β_3 is a subunit of the LRI (leucocyte responsive integrin), and is thought to be involved in inflammatory responses. Since all of the β_3 in the mutant mouse contains the tyrosine to phenylalanine mutations in the cytoplasmic domain, this mouse can be used to examine the effect of these mutations on $\alpha_v\beta_3$ and LRI function. Again, such analyses
15 will provide information on whether or not inhibition of β_3 phosphorylation will have effects on angiogenesis, tumor metastasis and/or inflammation, and thus will be of significant clinical import.

The foregoing detailed description has been given for clearness of understanding only and no unnecessary limitations should be understood therefrom as modifications will
20 be obvious to those skilled in the art.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present
25 disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth and as follows in the scope of the appended claims.

II. Specific Embodiments

30 The presence of phosphotyrosine binding motifs within the cytoplasmic domain of β_3 suggests that tyrosine phosphorylation of this integrin tail facilitates the recruitment of

phosphotyrosine-binding signaling proteins *Shc* and *Grb2* to the cell membrane. See Philips *et al.* (1996). The present inventors have discovered that tyrosine phosphorylation and dephosphorylation of the $\beta 3$ integrin tail may, in an analogous manner, regulate integrin association with the cytoskeleton (see, e.g., U.S. Patent Application No.

5 08/975,653, filed November 21, 1997, which is incorporated by reference herein). Figure 2 provides the amino acid sequences of various human integrin β subunits, showing the locations of the tyrosine residues. As mentioned previously, other studies have shown that the sequences of the cytoplasmic domains of $\beta 1$, $\beta 2$ and $\beta 3$ (GP IIIa) which contain tyrosines are important for normal functioning of GPIIb-IIIa and of other integrins. Thus, 10 transgenic animals that express altered and mutant cytoplasmic domains of those or other integrin β subunits that natively contain phosphorylatable tyrosine residues are expressly included in the scope of this invention.

Also expressly contemplated are altered, modified or mutated integrin subunits that contain conservative substitutions for the phosphorylatable tyrosine residues. As used 15 herein, a conservative amino acid substitution refers to alterations of particular residues (specifically the phosphorylatable tyrosine residues) or other alterations in the amino acid sequence that, except for the inability to undergo phosphorylation at the site of the one or more relevant cytoplasmic tyrosine residues, do not otherwise adversely affect the ability of the integrin to participate in signaling activities or to bind to one or more particular 20 binding partners, such as myosin, for example. A substitution, insertion or deletion is said to adversely affect the normal functioning of the corresponding wild-type integrin β subunit peptide when the altered sequence significantly inhibits the peptide from participating in signalling activities or from associating with its native binding partners, such as myosin. For example, it is contemplated that the overall charge, structure or 25 hydrophobic/hydrophilic properties of the β subunit peptide can be altered without adversely affecting activity of the peptide. Accordingly, it is contemplated that the amino acid sequence of the β subunit peptide can be altered, for example to render the peptide more hydrophobic or hydrophilic, to the extent that this does not adversely affecting the ability of the peptide to otherwise participate in its normal signalling

activities or to become associated with normal binding partners, such as myosin. Thus, specifically contemplated are mutant GPIIIa subunits which are

For example, in one application, altered β subunit peptides can be generated using standard knock-out procedures to modify any one or all of the phosphorylatable tyrosine residues. This can be accomplished using a variety of art-known procedures such as targeted recombination. Once generated, the genetically-engineered animal can be used to

- 1) identify biological and pathological processes mediated by the β subunit peptide;
- 2) identify proteins and other genes that interact with the β subunit peptide; 3) identify agents that can be exogenously supplied to overcome the absence or reduction in phosphorylation of the β subunit peptide; and 4) serve as an appropriate screen for identifying agents that modulate (*i.e.*, increase or decrease) the activity of the altered β subunit peptides.

As discussed herein, the present invention provides transgenic animals that contain non-tyrosine residues at sites where phosphorylatable tyrosine residues occur normally in endogenous β subunit genes. Since the non-tyrosine substitutions will not be phosphorylated as in the case where the residues are tyrosine and since normal platelet aggregation is dependent on phosphorylation occurring, the transgenic mammals of the present invention will display non-normal platelet aggregation. By comparing the physiological and morphological characteristics between the transformed and non-transformed animals one skilled in the art can thereby determine the effect of the presence or absence of cytoplasmic tyrosine phosphorylation in the GP IIIa gene on the corresponding animal.

Furthermore, the transgenic animals of the present invention can be used to identify agents that modulate (*i.e.*, either promote or further inhibit) platelet aggregation or other effects that are mediated by integrin signaling pathways. The evaluation of such agents can be conducted either *in vitro*, *in situ*, or *in vivo* by techniques known to those skilled in the art. The cells, platelets, tissues and whole organisms of the disclosed transgenic animals have utility in testing the effect of various agents for their ability to reduce or increase the processes involved with integrin-mediated cytoskeletal association. Agents which can be tested include various anticoagulant, thrombolytic and antiplatelet therapeutics and drugs. Examples of such agents include glycosaminoglycans such as

heparin; oral anticoagulants such as dicumarol, anisindione, and bromadkiolone; tissue plasminogen activator (t-PA); urokinase; aspirin; dipyridamole; and ticlopidine. See, Majerus, et al., *Anticoagulant, Thrombolytic, and Antiplatelet Drugs*, in Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, Ninth Edition, Chapter 54 (1996)

5 for a more complete list of such agents and their pharmacology.

The cells and whole organisms of the transgenic animals of the present invention, quite apart from their uses in veterinary and human medicine, may also be used to investigate gene regulation, expression and organization in animals. By studying the various effects of altered or mutant integrin β subunits, for example the GP IIIa gene, on
10 other cells, tissues and biological processes, one skilled in the art can ascertain the role of tyrosine phosphorylation in the normal metabolic pathways.

For further examples of diagnostic and research uses of transgenic mammals, especially transgenic mice, see U.S. Patent No. 5,569,824.

Without further description, it is believed that one of ordinary skill in the art can,
15 using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. The following working examples therefore, specifically point out preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

20 Other generic configurations will be apparent to one skilled in the art.

Example 1

Obtaining murine genomic GP IIIa DNA

The sequence for the murine genomic DNA is not known and has not been published,
25 however part of the amino acid sequence of mouse GP IIIa was available (Cietat et al. (1993) *Biochem et Biophys Res Comm.* 193: 771-778, and Dr Jean-Phillipe Rosa, *Unite INSERM* 348, Paris) and its similarity to human GP IIIa sequence suggested the genomic GP IIIa from humans and mice could be fairly similar. Therefore, several PCR primers were generated towards the mouse GP IIIa sequence in areas which, in the case of human GP IIIa, spanned the

murine genomic DNA and one set generated a specific band of approximately 1.5kB in the PCR (polymerase chain reaction). This band was close in size to the predicted human fragment (~1.2kB) which spanned the 3' part of exon M, the 5' part of exon N, and the intron between these two exons. The primers encoded: 1) 5' primer - residues 1834-1850 and 2) 3' primer - residues 2026-2039 of murine GP IIIa. These specific primers were then used to obtain a P1 clone containing a large fragment of murine genomic DNA which gave the same band when used as a template in PCR. Thus, we had obtained a P1 clone that contained murine genomic GP IIIa, and specifically contained the regions encoding the cytoplasmic domain of GP IIIa. Next, ~10kB of this P1 clone was mapped by a mixture of restriction digests, Southern Blotting and amino acid sequencing. It was confirmed that this was murine GP IIIa and that, similar to human GP IIIa, two exons (termed M and N after the human nomenclature) encoded the murine GP IIIa cytoplasmic region. Indeed the cytoplasmic domain amino acid residues turned out to be identical between human and mouse.

Example 2

Mutation of tyrosine residues 747 and 759 to phenylalanines

A .3kB PstI/SacI fragment containing exon N (which encodes the two tyrosine residues) was subcloned into pBSKS II and the two tyrosine residues were mutated to phenylalanines using standard site-directed mutagenesis techniques. Mutations were confirmed by sequencing and were also designed to introduce two new enzyme sites, a DraI site (at Y747F) and a TaqI site (at Y759F) which could be used in later identification of the mutant DNA. This mutant fragment was used to replace the wild-type PstI/SacI fragment in the murine genomic GP IIIa clone.

Example 3

Construction of targeting vector

An approximately 2.5kB EcoRI/HindIII fragment of murine GP IIIa, containing exon M as well as intronic genomic DNA was subcloned, using HindIII/EcoRI adaptors, into an EcoRI site of a pBS vector containing a neomycin resistant cassette. The neo^r DNA was flanked by FRT recognition sequences which are sites recognized by the Flp recombinase that

can be used to excise the neo^r DNA at a later stage (Dymecki S. M. (1996) *PNAS USA* 93: 6191-6196). A ~4kB HindIII fragment of GP IIIa containing exon N with the tyrosine to phenylalanine mutations was subcloned into a HindIII site on this vector to give a construct basically consisting of the mutated GP IIIa with a neo^r cassette in the intron between exons M and N.

Example 4

Generation of targeted embryonic stem (ES) cells

The above targeting construct was transfected into ES cells which were then selected for neo^r using the drug G418 as described in Johnson, R and Killeen, N. P (*In Gene Probes 2: a practical approach*, Eds. B. D. Hanes and S. J. Higgins. Oxford University Press p. 313-327 (1995)). ES cell which had undergone a successful homologous recombination event were identified by Southern blotting and PCR techniques. These cells now had one wild type GP IIIa allele and one containing the neo^r cassette and the tyrosine to phenylalanine mutations. The drug cassette is 1.9kB in size and although it was present in an intron we wished to remove it to make sure that its presence would have no detrimental effect in our experiments. To this effect the selected ES cells were transiently transfected with the Flp recombinase which recognized the FRT sites and led to the excision of the DNA encoding the drug resistance. Thus, instead of a 1.9kB piece of extraneous DNA being present in the intron a small piece encoding the 34bp FRT recognition site (and a new XbaI site) is all that remains (Dymecki, S. M., *PNAS* 93: 6191-6196 (1996)). Again, ES cells that had undergone a successful event were selected for using Southern blotting and PCR techniques.

Example 5

Generation of mutant mice

To produce mutant mice a mutant ES cell clone (originally from a strain 129 mouse) was injected into blastocysts from a C57Bl6 mouse and these blastocysts were implanted into a pseudo-pregnant foster mother (as in Ramirez-Solis, R., Davis, A. C., and Bradley, A. *Guide to techniques in mouse development*. Methods in Enzymology 225. Eds. P. M. Wassarman

ES cells) were then mated with C57Bl6 wild-type females. The offspring were genotyped for the presence of wildtype and mutant GP IIIa alleles by Southern blotting (making use of the new XbaI site that was present only in mutant GP IIIa DNA). The offspring were a mix of wild-type and heterozygote animals and the heterozygote animals were further crossed to
5 produce litters containing wildtype animals, heterozygote animals (ie. one wildtype and one mutant GP IIIa allele) and homozygotes (ie. both GP IIIa alleles containing the tyrosine to phenylalanine mutations).

To date the genotypes of the offspring from such matings have occurred in the expected ratio (assuming basic Mendelian genetics). Furthermore, the mutant animals are
10 viable and express GP IIb-IIIa on their platelets at similar levels to that seen in normal animals expressing non-mutant GP IIIa when the platelets are stained with an anti-GP IIb-IIIa antibody and examined on the FACS (fluorescent activated cell sorter).

Although the present invention has been described in detail with reference to examples above, it is understood that various modifications can be made without departing from the
15 spirit of the invention. Accordingly, the invention is limited only by the following claims. All cited patents, applications and publications referred to in the application are hereby incorporated by reference in their entirety.

WHAT IS CLAIMED:

1. A non-human mammal comprising a mutant GP IIIa gene wherein at least one of the two cytoplasmic tyrosine residues of the gene has been replaced with a non-tyrosine
5 residue.

2. The non-human mammal of claim 1 wherein the non-tyrosine residue is phenylalanine.

10 3. The non-human mammal of claim 1 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759.

4. Platelets isolated from the blood plasma of the non-human mammal of claim 1.

15 5. The non-human mammal of claim 1 wherein the non-human mammal is selected from the group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

20 6. The non-human mammal of claim 5 wherein the non-human mammal is a mouse.

7. The non-human mammal of claim 1 wherein both cytoplasmic tyrosine residues have been replaced with a non-tyrosine residue.

25 8. The non-human mammal of claim 7 wherein the non-tyrosine residues are phenylalanine.

9. The non-human mammal of claim 7 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759

11. The non-human mammal of claim 7 wherein the non-human mammal is selected from the group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

5

12. The non-human mammal of claim 11 wherein the non-human mammal is a mouse.

13. A non-human mammal expressing a transgene stably introduced into its DNA, wherein the transgene comprises DNA encoding mutant GP IIIa wherein at least one of the two cytoplasmic tyrosine residues has been replaced with a non-tyrosine residue.

10

14. The non-human mammal of claim 13 wherein the non-tyrosine residue is phenylalanine.

15

15. The non-human mammal of claim 13 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759.

16. Platelets isolated from the blood plasma of the non-human mammal of claim 13.

20

17. The non-human mammal of claim 13 wherein the non-human mammal is selected from the group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

25

18. The non-human mammal of claim 17 wherein the non-human mammal is a mouse.

19. The non-human mammal of claim 13 wherein both cytoplasmic tyrosine residues have been replaced with a non-tyrosine residue.

30

20. The non-human mammal of claim 19 wherein the non-tyrosine residues are phenylalanine.

21. The non-human mammal of claim 19 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759.

22. Platelets isolated from the blood plasma of the non-human mammal of claim 19.

23. The non-human mammal of claim 19 wherein the non-human mammal is selected from the group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

24. The non-human mammal of claim 23 wherein the non-human mammal is a mouse.

25. A method of preparing a transformed non-human mammal with a mutant GP IIIa gene wherein at least one of the two tyrosine residues of the endogenous GP IIIa gene has been replaced with a non-tyrosine residue to prepare the mutant GP IIIa, said method comprising:

- a) introducing into embryonic stem cells a nucleic acid molecule encoding the mutant GP IIIa gene;
- b) regenerating a transformed non-human mammal from the cells resulting from step a).

26. The method of claim 25 wherein the non-tyrosine residue is phenylalanine.

27. The method of claim 25 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759.

28. The method of claim 25 wherein the non-human mammal is selected from the group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

5 29. The method of claim 28 wherein the non-human mammal is a mouse.

30. The method of claim 25 wherein both cytoplasmic tyrosine residues have been replaced with a non-tyrosine residue.

10 31. The method of claim 30 wherein the non-tyrosine residues are phenylalanine.

32. The method of claim 30 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759.

15 33. The method of claim 30 wherein the non-human mammal is selected from the group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

20 34. The method of claim 33 wherein the non-human mammal is a mouse.

35. The method of claim 25 further comprising breeding the transformed non-human mammal so as to produce a non-human mammal homozygotic for the mutant GP IIIa gene.

25 36. The method of claim 35 wherein the non-human mammal is a mouse.

37. A method of preparing a transformed non-human mammal with a mutant GP IIIa gene wherein at least one of the two tyrosine residues of the endogenous GP IIIa gene has been replaced with a non-tyrosine residue to prepare the mutant GP IIIa, said method
30 comprising:

- 5
- a) introducing into embryonic stem cells a nucleic acid molecule encoding the mutant GP IIIa gene and a selectable marker flanked by FRT sites;
 - b) identifying and selecting transformed cells;
 - c) removing the selectable marker from the transformed cells selected in step b) by transient transformation with FLP recombinase;
 - d) injecting the transformed cells from step c) into blastocysts; and,
 - e) regenerating a transformed non-human mammal from the blastocysts of step d), wherein the regenerated transformed non-human mammal is chimeric for the mutant GP IIIa gene.
- 10

38. The method of claim 37 wherein the non-tyrosine residue is phenylalanine.

39. The method of claim 37 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759.

15

40. The method of claim 37 wherein the non-human mammal is selected from the group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

20

41. The method of claim 37 wherein the non-human mammal is a mouse.

42. The method of claim 37 wherein both cytoplasmic tyrosine residues have been replaced with a non-tyrosine residue.

25

43. The method of claim 37 wherein the non-tyrosine residues are phenylalanine.

44. The method of claim 37 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759.

45. The method of claim 43 wherein the non-human mammal is selected from the group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

5 46. The method of claim 45 wherein the non-human mammal is a mouse.

47. The method of claim 37 further comprising breeding the transformed non-human mammal so as to produce a non-human mammal homozygotic for the mutant GP IIIa gene.

10

48. The method of claim 47 wherein the non-human mammal is a mouse.

49. The method of claim 37 further comprising the following steps:

15

f) breeding the chimeric non-human mammal with a wild-type non-human mammal to produce a non-human mammal heterozygotic for the mutant GP IIIa gene;

g) crossing a heterozygotic non-human mammal produced in step f) with a second heterozygotic non-human mammal produced in step f) ; and,

20

h) selecting a non-human mammal homozygotic for the mutant GP IIIa gene from the resulting progeny.

50. The method of claim 49 wherein the non-human mammal is a mouse.

25

51. A method comprising comparing a characteristic between two mammals of the same species, wherein one mammal has a wild-type GP IIIa gene and the other mammal has a mutant GP IIIa gene, wherein at least one of the two tyrosine residues of the wild-type GP IIIa gene has been replaced with a non-tyrosine residue in the mutant GP IIIa gene.

30

52. The method of claim 51 wherein the non-tyrosine residue is phenylalanine.

53. The method of claim 51 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759.

54. The method of claim 51 wherein the non-human mammal is selected from the
5 group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat, rabbit, cow and guinea pig.

55. The method of claim 54 wherein the non-human mammal is a mouse.

10 56. The method of claim 51 wherein both cytoplasmic tyrosine residues have been replaced with a non-tyrosine residue.

57. The method of claim 56 wherein the non-tyrosine residues are phenylalanine.

15 58. The method of claim 56 wherein the cytoplasmic tyrosine residues are tyrosine residues 747 and 759.

59. The method of claim 56 wherein the non-human mammal is selected from the
group consisting of sheep, goat, mouse, pig, dog, cat, monkey, chimpanzee, hamster, rat,
20 rabbit, cow and guinea pig.

60. The method of claim 59 wherein the non-human mammal is a mouse.

61. The method of claim 51 further comprising comparing the bleeding time
25 between the two mammal types.

62. The method of claim 51 further comprising comparing the thrombotic responses between the two mammal types.

64. The method of claim 51 further comprising comparing tumor metastasis between the two mammal types.

5 65. The method of claim 51 further comprising comparing inflammation between the two mammal types.

66. The method of claim 51 wherein the mammal is a mouse.

10 67. A method of determining the effect of an agent on a characteristic of a mammal that is attributable to the expression of the GP IIIa gene, said method comprising;

- a) administering said agent to the mammal of claim 1;
- b) maintaining said mammal for a desired period of time after said administration; and,
- 15 c) determining whether a characteristic of said mammal that is attributable to the expression of the mutant GP IIIa gene has been affected by the administration of said agent.

20 68. The method of claim 67 wherein the mammal is a mouse.




FIG. 1

GP III a (β3)	KLLLTTHDRK	EFAKFEeEra	rAkWdtanNP	LYKeAt ⁵ IftNi	tYrgt.....
β6	KLLVsfHDRK	EvAKFEaErs	kAkWqtgNP	LYrgststfkNv	tYfhreKqkv dlstdc
β1	KLLmIHDRR	EEAKEEEKEm	rAkWdtgNP	iYKsAvtIvvNp	kYegk.....
β5	KLLVtIHDRR	EFAKFqSErs	rArYemasNP	LYrkpiStht	vdftfnkfnk	sYngtvd....
β2	KaLThIsDIR	EYrrFEkEkI	ksqWnnd.NP	LFKsAltIvmNp	kFaes.....
β7	rLsVeIyDRR	EYsrFEkEqq	qInwkqdsNP	LYKsAltItiNp	rFqeadsptI
Cn	KLLV-IHDRR	EFAKFE-E--	-A-W----	NP LYK-A--T--	-----N-	-Y-----

FIG. 2

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/08285

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N5/10 A01K67/027 C07K14/705 A61K49/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation - first classification system followed by classification system

IPC 6 C12N A01K C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>BLYSTONE S ET AL: "Requirement of integrin beta3 tyrosine 747 for beta3 tyrosine phosphorylation and regulation of alphavbeta3 avidity" JOURNAL OF BIOLOGICAL CHEMISTRY.. vol. 272, no. 45, 7 November 1997 (1997-11-07), pages 28757-28761, XP002112506 figure 2</p> <p>---</p> <p>-/--</p>	1-68

☒ Further documents are listed in the continuation of this report

☒ Patent family members are listed in annex

Special categories of cited documents

A: document defining the general state of the art which is not considered to be of particular relevance

E: earlier document but published on or after the international filing date

L: document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

R: document relevant to the prior art of the invention

* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other documents

18 August 1999

18 August 1999

Name of the applicant

Name of the applicant

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/08285

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate of the relevant passages	Relevant to claim No
Y	<p>LIU K ET AL: "Identification of a functionally important sequence in the cytoplasmic tail of integrin beta 3 by using cell-permeable peptide analogs" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA., vol. 93, no. 21, 15 October 1996 (1996-10-15), pages 11819-11824, XP002112507 cited in the application figure 6</p>	1-68
Y	<p>HANKS M ET AL: "Rescue of the En-1 mutant phenotype by replacement of En-1 with En-2" SCIENCE., vol. 269, no. 5224, 4 August 1995 (1995-08-04), pages 679-682, XP002112508 figure 1A</p>	1-68
Y	<p>BAUDOIN C ET AL: "Knockout and knockin of the betal exon D define distinct roles for integrin splice variants in heart function and embryonic development" GENES & DEVELOPMENT., vol. 12, no. 8, 15 April 1998 (1998-04-15), pages 1202-1216, XP002112509 figures 1,4</p>	1-68
A	<p>WO 97 08316 A (MERCK & CO INC ; DUONG LE T (US); NUTT ELKA M (US); RODAN GIDEON A) 6 March 1997 (1997-03-06)</p>	
A	<p>WO 95 25173 A (SCRIPPS RESEARCH INST) 21 September 1995 (1995-09-21)</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/08285

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9708316 A	06-03-1997	CA 2229622 A	06-03-1997
		EP 0854920 A	29-07-1998
		US 5895754 A	20-04-1999
W0 9525173 A	21-09-1995	US 5523209 A	04-06-1996
		CA 2185446 A	21-09-1995
		EP 0769065 A	23-04-1997
		JP 9510609 T	28-10-1997

